

Scotland's Rural College

## Results of an international phosphorus digestibility ring test with broiler chickens

Rodehutschord, M; Adeola, O; Angel, R; Bikker, P; Delezie, E; Dozier III, WA; Umar Faruk, M; Francesch, M; Kwakernaak, C; Narcy, A; Nyachoti, CM; Olukosi, OA; Preynat, A; Renouf, B; Saiz del Barrio, A; Schedle, K; Siegert, W; Steinfeldt, S; van Krimpen, MM; Waititu, SM; Witzig, M

*Published in:*  
Poultry Science

*DOI:*  
[10.3382/ps/pew426](https://doi.org/10.3382/ps/pew426)

First published: 05/12/2016

*Document Version*  
Peer reviewed version

[Link to publication](#)

### *Citation for pulished version (APA):*

Rodehutschord, M., Adeola, O., Angel, R., Bikker, P., Delezie, E., Dozier III, WA., Umar Faruk, M., Francesch, M., Kwakernaak, C., Narcy, A., Nyachoti, CM., Olukosi, OA., Preynat, A., Renouf, B., Saiz del Barrio, A., Schedle, K., Siegert, W., Steinfeldt, S., van Krimpen, MM., ... Witzig, M. (2016). Results of an international phosphorus digestibility ring test with broiler chickens. *Poultry Science*, 96(6), 1679 - 1687.  
<https://doi.org/10.3382/ps/pew426>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 05. May. 2023

## PHOSPHORUS DIGESTIBILITY

### Results of an international phosphorus digestibility ring test with broiler chickens

M. Rodehutsord,<sup>\*1</sup> O. Adeola,<sup>†</sup> R. Angel,<sup>‡</sup> P. Bikker,<sup>∞</sup> E. Delezie,<sup>=</sup> W. A. Dozier III,<sup>§</sup>  
M. Umar Faruk,<sup>#</sup> M. Francesch,<sup>||</sup> C. Kwakernaak,<sup>l</sup> A. Narcy,<sup>♦</sup> C. M. Nyachoti,<sup>▲</sup>  
O. A. Olukosi,<sup>¥</sup> A. Preynat,<sup>\$</sup> B. Renouf,<sup>~</sup> A. Saiz del Barrio,<sup>Ψ</sup> K. Schedle,<sup>¶</sup> W. Siegert,<sup>\*</sup> S.  
Steenfeldt,<sup>&</sup> M. M. van Krimpen,<sup>∞</sup> S. M. Waititu,<sup>▲</sup> and M. Witzig<sup>\*</sup>

\*Institute of Animal Science, University of Hohenheim, 70599 Stuttgart, Germany

†Department of Animal Sciences, Purdue University, West Lafayette, IN 47907

‡Department of Animal and Avian Sciences, University of Maryland, College Park, MD 20742

∞Wageningen UR Livestock Research, 6700 AH Wageningen, The Netherlands

=Animal Sciences Unit, Institute for Agricultural and Fisheries Research (ILVO), 9090 Melle, Belgium

§Department of Poultry Science, Auburn University, Auburn, AL 36849

#DSM Nutritional Products, 68128 Village-Neuf, France

||IRTA, Mas de Bover, 43120 Constantí (Tarragona), Spain

lSchothorst Feed Research BV, 8200 AM Lelystad, the Netherlands

♦INRA-UR0083, Recherches Avicoles, 37380 Nouzilly, France

▲Department of Animal Science, University of Manitoba, Winnipeg, MB, Canada R3T 2N2

¥Monogastric Science Research Centre, Scotland's Rural College, Edinburgh, EH9 3JG, UK

\$Adisseo – Cern, 6 route noire, 03600 Malicorne, France

~SAS Euronutrition, Domaine Expérimental le Pavillon, 72240 Saint Symphorien, France

ΨTrouw Nutrition R&D Poultry Research Centre, 45950 Casarrubios del Monte, Spain

¶Institute of Animal Nutrition, Livestock Products and Nutrition Physiology, University of  
Natural Resources and Life Sciences, Vienna, 1190 Vienna, Austria

&Department of Animal Science, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark

### Scientific Section: Metabolism and Nutrition

<sup>1</sup>Corresponding author: inst450@uni-hohenheim.de

Markus Rodehutsord, University of Hohenheim, Institute of  
Animal Science, Emil-Wolff-Straße 10, 70599 Stuttgart, Germany

**ABSTRACT** The objective of this ring test was to investigate the prececal phosphorus (P) digestibility of soybean meal (SBM) in broiler chickens using the trial protocol proposed by the World's Poultry Science Association. It was hypothesized that prececal P digestibility of SBM determined in the collaborating stations is similar. Three diets with different inclusion levels of SBM were mixed in a feed mill specialized in experimental diets, and transported to 17 collaborating stations. Broiler chicks were raised on commercial starter diets according to station-specific management routine. Then they were fed the experimental diets for a minimum of 5 d before content of the posterior half of the ileum was collected. A minimum of 6 experimental replicates per diet was used in each station. All diets and digesta samples were analyzed in the same laboratory. Diet, station, and their interaction significantly affected ( $P < 0.05$ ) the prececal digestibility values of P and calcium of the diets. The prececal P digestibility of SBM was determined by linear regression and varied between stations from 19 to 51 %, with significant differences among stations. In a subset of 4 stations, the prececal disappearance of *myo*-inositol 1,2,3,4,5,6-hexakis (dihydrogen phosphate)-P;  $\text{InsP}_6\text{-P}$  was also studied. The prececal  $\text{InsP}_6\text{-P}$  disappearance correlated well with the prececal P digestibility. We hypothesized that factors influencing  $\text{InsP}_6$  hydrolysis were main contributors to the variation in prececal P digestibility between stations. These factors were probably related to the feeding and housing conditions (floor pens or cages) of the birds in the pre-experimental phase. Therefore, we suggest that the World's Poultry Science Association protocol for the determination of digestible P be extended to the standardization of the pre-experimental period. We also suggest that comparisons of P digestibility measurements between studies are only made with great caution until the protocol is more refined.

**Key words:** broiler chickens, phosphorus, phytate, prececal digestibility, variability

## INTRODUCTION

Phosphorus (P) is an element of high relevance for poultry feeding. Diets are usually supplemented with feed phosphates or phytase or both in order to fulfil the animal's requirement for available P. However, excessive intake can contribute to environmental problems in areas with high livestock density. The variation in P availability of different feed raw materials is high (Shastak and Rodehutscord, 2015), and it is generally accepted that the use of P as a globally finite resource can be optimized by considering the differences that exist in P availability of feed raw materials.

Different response criteria and descriptive terms for available P have been used in the literature of the past seven decades (Shastak and Rodehutscord, 2013). These differences make it difficult to compare results obtained by using different techniques in different laboratories, and to compile comprehensive feedstuff tables needed by the industry. In an attempt to improve this situation the Working Group No 2: Nutrition of the European Federation of Branches of the World's Poultry Science Association proposed a standard protocol for the determination of P availability (WPSA, 2013). This protocol is based on using the digestibility measured at the terminal ileum of broiler chickens (prececal digestibility of P (**pcdP**), otherwise also referred to as ileal digestibility). The protocol defines assay details relevant for the outcome of the measurement, such as age of birds, minimum number of experimental replicates, diet composition, and P and calcium (Ca) levels in the diet.

The expectation from implementing this standard protocol is that results generated in different research stations for the same feed raw material are similar and thus can better be compared and used by the industry. However, application of the standard protocol has not been compared yet between stations. Thus, our objective was to compare the results between our laboratories for one feed raw material when based on the WPSA (2013) standard protocol.

We choose soybean meal (**SBM**) for this comparison, and we hypothesized that the pcdP of SBM determined in our laboratories is similar.

## MATERIALS AND METHODS

Seventeen research stations from Europe and North America collaborated in this study and determined the pcdP of SBM following the principles of the standard protocol of WPSA (2013), which includes regression analysis. The concept of this ring test required that all stations use the same diets and thus variability in raw material quality and diet preparation was mitigated. Furthermore, chemical analyses of all diets and ileal digesta collected in all stations were conducted in the same laboratory. Customs regulations related to cross-continental shipment of feed samples made it impossible for more laboratories from other regions of the world to participate in the ring test.

### *Experimental diets*

Three diets that differed in the inclusion level of SBM (diets A, B, C) were formulated based on the examples of WPSA (2013) (Table 1). The de-hulled, solvent-extracted SBM was included at the expense of corn starch, thus making SBM the only source of variation in P content of the diets. Analytical characterization of the SBM is presented as a footnote in Table 1. Titanium dioxide was included (0.5 %) as the indigestible marker. The concentrations of total P and phytic acid (*myo*-inositol 1,2,3,4,5,6-hexakis (dihydrogen phosphate)-P; **InsP<sub>6</sub>-P**) in the diets ranged from 3.0 to 4.6 and 1.6 to 2.3 g/kg dry matter (DM), respectively. The Ca:total P ratio was analyzed in the diets to be 1.5:1.0, which was very close to the ratio recommended by WPSA (2013) (1.3:1.0 - 1.4:1.0). Diets were manufactured at Research Diet Services (location Wijk bij Duurstede, Netherlands). First, all ingredients except the variable ones (SBM, corn starch and limestone) were mixed in one lot. Then, diets A and C were mixed by adding the respective amounts of SBM, corn starch and limestone. Finally, diet B

was mixed by 50 % diet A and 50 % diet C. All diets were pelleted (die hole: 3.0 mm x 17 mm) with a little steam addition. Temperature of the pellets measured directly after leaving the die was 77°C for diets A and B and 74°C for diet C. The experimental diets were bagged, sealed, and transported to the 17 participating stations.

### ***Birds and experimental procedures***

Animal trials at the participating stations were conducted between March and September 2014. All animal procedures were in accordance with the animal welfare regulations that were applicable for the individual participating station. While basic principles of the WPSA (2013) protocol were applied in all laboratories, details of the trials were specific for each station (Table 2). Broiler chicks (Ross 308, Ross 708, Ross PM3, and Heritage 5632) were fed commercial starter feed until they were between 12 and 23 d old and when the treatment diets were introduced. At least 6 experimental replicates were used per diet by each station, and each replicate had a minimum of 8 birds. One laboratory used only 4 birds per replicate but 10 replicates per diet. Feed and drinking water were offered for ad libitum consumption. The ADFI and average BW were recorded. Birds were fed the experimental diets for a minimum period of 5 d and then were sacrificed at an age of 21-28 d. The abdominal cavity was immediately opened, the digestive tract removed, and the ileum (section between Meckel's diverticulum and 2 cm anterior to the ileo-ceco-colonic junction) dissected. The digesta of the distal half of the ileum was obtained by flushing with water or by gentle squeezing. Digesta from all birds of one replicate were pooled into one sample. Samples were dried (freeze-drying or oven drying), ground, and sent to the Hohenheim laboratory together with the respective diet samples for chemical analyses.

### ***Chemical Analyses***

Proximate nutrients in the SBM and CP concentration in the diets were analyzed according to the official methods (Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten (VDLUFA, 2007)). Analysis of Ca, P, and Ti in diets and digesta samples was performed using an inductively coupled plasma optical emission spectrometer following a sulfuric and nitric acid wet digestion with all specifications described by Zeller et al. (2015b). Concentrations of InsP<sub>6</sub> and lower inositol phosphates in the diets were analyzed following EDTA extraction at pH 10 using high-performance ion chromatography as described by Zeller et al. (2015c). Digesta samples from 4 stations that had freeze-dried the samples and were found to be different in pcdP values were also analyzed for InsP<sub>6</sub> using the same method as used for the diets. This was not planned from the beginning and is not part of the WPSA (2013) protocol. Hence, some stations oven-dried the samples. Oven dried samples were not considered for InsP<sub>6</sub> analysis because some InsP<sub>6</sub> might be hydrolyzed after sampling from the ileum during the drying process.

### *Calculations and Statistics*

Prececal digestibility of P and Ca, and disappearance of InsP<sub>6</sub> (y) was calculated on a pen basis according to the following equation:

$$y (\%) = 100 - 100 \times \left( \frac{\text{Ti in the diet (g/kg DM)}}{\text{Ti in the digesta (g/kg DM)}} \right) \times \left( \frac{\text{InsP}_6 \text{ or P or Ca in the digesta (g/kg DM)}}{\text{InsP}_6 \text{ or P or Ca in the diet (g/kg DM)}} \right)$$

The amount of pcdP (g/kg DM) was calculated by multiplication of the P concentration of the diet (g/kg DM) and the respective digestibility (%) divided by 100.

Statistical evaluation of the data used the MIXED procedure of SAS for Windows (Version 9.3, SAS Institute, Cary, NC). Evaluation of P digestibility, Ca digestibility, and InsP<sub>6</sub> disappearance values of the diets was performed using diet (A, B, C), stations (1 - 16), and their interaction as fixed effects. The trial in one station generated questionable digestibility data for P and Ca that could not be explained. Hence data of this trial were not included in the data evaluation and only 16 trials were used.

Digestibility of P from SBM was calculated for each of the 16 stations by linear regression. Linear regressions of the type  $y = a + mx$  were calculated using the SOLUTION statement to describe the relationship between pcdP content and P content in the diet (both in g/kg DM) for each station. Because differences in the P content between diets originated only from SBM inclusion, calculated slopes multiplied by 100 are regarded as an estimate of the pcdP of SBM (WPSA, 2013). The  $R^2$  and root MS error are reported as measures for the goodness of fit. Differences in pcdP of SMB between stations were compared using the ESTIMATE statement.

## RESULTS

The average pcdP of the diets across all stations for diets A, B, and C were 67, 61, and 55 %, respectively (Figure 1). Effects of diet, station, and their interaction were statistically significant ( $P < 0.05$ ). The ranges of pcdP values of the diets across stations were 55 to 82 % for diet A, 46 to 79 % for diet B, and 45 to 69 % for diet C (Table 3).

When the pcdP concentration of the diets was regressed against total P concentration, estimated slopes of the linear regressions ranged between stations from 0.19 to 0.51 (Table 4). Even when the two lowest and two highest slopes were disregarded, the slopes of the remaining stations still ranged from 0.22 to 0.42 with significant differences among stations.

The average prececal (pc) disappearance of  $\text{InsP}_6$  as studied in 4 stations varied from 58 to 74 %, 43 to 67 %, and 23 to 58 % for diets A, B, and C, respectively (Table 5), thus confirming the trends in differences seen in pcdP of the diets. Increments in dietary  $\text{InsP}_6$  with increasing SBM level of the diet were not related to an increase of pc  $\text{InsP}_6$  disappearance (Figure 2). However, across all diets and the 4 stations, a relationship between the pc disappearance of  $\text{InsP}_6$ -P and pcdP content became apparent, indicating that a very high proportion of the  $\text{InsP}_6$ -P that disappeared until the end of the ileum was absorbed (Figure 3).



The average prececal digestibility of Ca (pcdCa) of the diets across all stations was 57 % (diet A), 51 % (diet B), and 46 % (diet C), with a similar range between stations as found for pcdP (Figure 1). Effects of diet, station, and their interaction on pcdCa were statistically significant ( $P < 0.05$ ). Within the range studied the relationship between pcdP and pcdCa in the diets was linear and not significantly different between diets (Figure 4). On average, the concentration of pcdCa in the diets increased by 0.97 g/kg DM with each 1.00 g/kg DM increase in pcdP.

## DISCUSSION

The underlying hypothesis of this study was that digestibility values determined for the diets and for the SBM are similar among stations. This hypothesis must be rejected on the basis of the results. The pcdP for the same SBM ranged between 19 and 51 % among stations, which can be a frustrating fact from the viewpoint of feed producers aiming to use P digestibility data in their feed formulation matrix. It is of note that the opposite was found in a ring test on pcd of amino acids in broilers (Ravindran et al., submitted). There, pcd of amino acids of a corn-SBM-based diet was similar among the 5 collaborative stations, once important protocol details had been agreed on. Probably there is something specific on the targeted nutrient (amino acids vs. P and InsP<sub>6</sub>) that caused high variation in one ring test but not in the other. It cannot be ruled out that this difference is related to hormonal control of absorption known for Ca and P, but not for amino acids. However, for reasons subsequently mentioned other factors are seen more likely involved.

In the current ring test we used the same diets in all stations to mitigate variation potentially caused by raw material origin or diet processing. All diet and digesta samples were analyzed in the same laboratory, hence minimizing variation of results potentially caused by differences in analytical protocols. In spite of this standardization, the large differences have occurred. This variation is most likely associated with how the experiment was conducted and

birds managed in each station. While most assay details are standardized in the applied protocol (WPSA, 2013), some were not and will be discussed herein.

The starter diets used in the pre-experimental period met the bird's requirements, but were specific for each station and not of the same ingredient composition. Digestive capacity adaptation to Ca and P deficiencies or imbalances can occur within 48 h (Angel et al., 2013). Hence, any adaptation to the experimental diets would have already occurred in the present study before samples were taken. However, it cannot be ruled out that starter diet details have affected the measurements subsequently made with the experimental diets. The P concentration of the starter diets (formulated values) was between 0.5 and 0.7 %, and the Ca concentration between 0.7 and 1.1 %. Some of the starter diets contained an added phytase while others did not. Compensatory adaptation in growth and bone mineralization in a later growth phase occurred when broilers were fed a diet moderately deficient in P and Ca from hatching to 18 d (Yan et al., 2005). A recent study showed that a low level of P in the diet fed from 10 to 21 d can affect the mRNA levels of several genes encoding Ca and P transporters at 35 d of age, depending on the amount of P and Ca fed from 22 to 35 d (Rousseau et al., 2016). These authors concluded that chickens are able to adapt to early dietary changes in P and Ca through improvement of digestive efficiency in a later phase. Starter diet P and Ca levels were different across participating stations in the current study, but none was deficient. It cannot be ruled out but not substantiated with the data that the differences affected the results. A negative trend might be interpreted when the P digestibility data are related to the P level of the starter diets (Figure 5A); however, the regression line did not significantly deviate from zero ( $P = 0.08$ ). A relationship with the formulated phytase content of the starter diet also was not apparent (Figure 5B). Because only the formulated values of P, Ca, and phytase are available for the starter diets, and because we do not have information about raw material composition of the starter diets, relationships between starter diet characteristics and determined digestibility values should be viewed with caution. In such complex design,

colinearity also between other factors of variation might occur (starter diet, duration of period, age, etc.) and thus could bias interpretation.

Some starter diets contained a coccidiostat, while others did not. As a consequence, microbiota colonization of the digestive tract might have developed differently between stations, which could affect InsP<sub>6</sub> hydrolysis as seen in the differences among stations and diets (Table 5). Previous studies have found a relatively high level of prececal or excreta InsP<sub>6</sub> disappearance in birds (25 to 78 %) even when diets devoid of any supplemental or plant-intrinsic phytase were used (Amerah et al., 2014; Applegate et al., 2003; Delezie et al., 2012; Li et al., 2016; Tamim et al., 2004; Zeller et al., 2015a; Zeller et al., 2015b). This range of values is high and probably related to dietary Ca levels or level and origin of InsP<sub>6</sub>. However, the generally high InsP<sub>6</sub> disappearance values strongly indicate that enzymes of non-feed origin, namely endogenous mucosal and bacterial phytases, have caused a substantial InsP<sub>6</sub> hydrolysis. Little is known so far about the contribution of specific bacteria to phytase activity in the digestive tract of chickens. However, lactobacilli were shown to be the main colonizers in the crop, jejunum, and ileum of broilers (Witzig et al., 2015). Some of the lactobacilli strains that were detected in gut content of broilers have been characterized to express high phytase activity (Palacios et al., 2008; Taheri et al., 2009). Use of the coccidiostat monensin reduced abundance of *Lactobacillus* sp. in ceca content (Danzeisen et al., 2011) and nisin-sensitive *L. reuteri* isolates were detected in the crop (Abbas Hilmi et al., 2007). Hence it is possible that coccidiostats included in some but not all starter diets used in the current ring test reduced abundance of bacteria known to possess phytase activity and thus InsP<sub>6</sub> hydrolysis. Initial microbiota colonization in the starter phase probably was effective in the experimental phase and contributed to InsP<sub>6</sub> hydrolysis from SBM to a different extent. We observed a very close relationship between InsP<sub>6</sub>-P disappearance and pcdP content of the diets (Figure 3), and although P of non-InsP<sub>6</sub> origin in SBM also contributed to the change in

pcdP content, the relationship well demonstrated that InsP<sub>6</sub> hydrolysis was a major determinant for the pcdP of SBM.

Another factor of potential relevance for microbial colonization and development of the digestive tract is the way birds were kept in the pre-experimental period. When birds were raised in floor pens, litter material intake could affect digestive tract development in general and microbial colonization in particular. Floor pen raising was practiced in some of the stations because required as per approval from the ethics committee or for other reasons.

Endogenous enzymes involved in intestinal InsP<sub>6</sub> hydrolysis can also originate from epithelial secretion because studies using purified brush border membrane vesicles from different sections of the small intestine of broiler chickens reported phytase activity (Huber et al., 2015; Maenz and Classen, 1998; Onyango et al., 2006). Significant differences in the  $V_{\max}$  of epithelial phytase kinetics were found between Hubbard  $\times$  Peterson and Ross  $\times$  Ross broilers, but not in pc InsP<sub>6</sub> hydrolysis (Applegate et al., 2003). It has been hypothesized from genetic studies that epithelial phytase expression is affected by the bird's genetic background (Beck et al., 2014). Indeed, genomic studies showed significant heritability in the range of 0.10 to 0.22 for P utilization, P excretion rate, and phytate-P bioavailability in broilers and Japanese quail (Beck et al., 2016; de Verdal et al., 2011; Zhang et al., 2003). In the current ring test, fast growing broilers from different strains were used that were typical for the respective station. It is possible that the genetic background of the parent lines used for broiler production in different regions of the world was different and this contributed to the differences we found among stations. However, it is not currently possible to quantify the contribution of epithelial phytase to intestinal luminal InsP<sub>6</sub> hydrolysis.

Some of the stations involved in the current study used unsexed broilers while others used males. As a side aspect of the current study one station has compared males and females when studying digestibility of the experimental diets (Schedle et al., 2016). The authors reported almost identical values and no significant differences between males and females.

This let us conclude that for the current study it was not relevant whether mixed-sex or male broilers were used.

Some other characteristics of the experiments such as growth, ADFI, age at sampling and killing procedures were also different among stations. Statistical search for any relationship would not be meaningful because the study was not designed to investigate any of the effects and the data set is too limited for correlation analysis. However, it does not appear from regressions that age at slaughter or at start of experimental feeding, BW at slaughter, and ADFI in the trial period affected the results (Figure 5).

The protocol of WPSA (2013) has standardized several details for the determination of P digestibility. From the results of this ring test, we conclude that the protocol standardization must go beyond the standards already set. Differences in starter diet composition and management conditions during the pre-experimental period of the trials affected the results and thus also need standardization. Until a more standardized protocol is established, care must be exercised when comparing P digestibility data from different laboratories.

## ACKNOWLEDGEMENTS

The financial support for this project provided by BASF SE, Nutreco, and Adisseo is gratefully acknowledged. Authors are also grateful to M. Schollenberger, M. Liebscher, and H. Ott for conducting chemical analyses.

## REFERENCES

Abbas Hilmi, H. T., A. Surakka, J. Apajalahti, and P. E. J. Saris. 2007. Identification of the most abundant *Lactobacillus* species in the crop of 1- and 5-week-old broiler chickens. *Appl. Environm. Microbiol.* 73:7867-7873. doi 10.1128/aem.01128-07

- 279 Amerah, A. M., P. W. Plumstead, L. P. Barnard, and A. Kumar. 2014. Effect of calcium level  
280 and phytase addition on ileal phytate degradation and amino acid digestibility of  
281 broilers fed corn-based diets. *Poult. Sci.* 93:906-915. doi 10.3382/ps.2013-03465
- 282 Angel, R., M. Proszkowiec-Weglarz, E. Jimenez-Moreno, S. W. Kim, and P. Plumstead.  
283 2013. Impact of time and dietary calcium and phosphorus deficiencies on their  
284 digestibilities in single ingredients. In: German branch of the WPSA (ed.) *Proc. 19th*  
285 *European Symposium on Poultry Nutrition*. Potsdam, Germany, August 26- 28, 2013,  
286 p. 199
- 287 Applegate, T. J., R. Angel, and H. L. Classen. 2003. Effect of dietary calcium, 25-  
288 hydroxycholecalciferol, or bird strain on small intestinal phytase activity in broiler  
289 chickens. *Poult. Sci.* 82:1140-1148.
- 290 Beck, P., H.-P. Piepho, M. Rodehutsord, and J. Bennewitz. 2016. Inferring relationships  
291 between phosphorus utilization, feed per gain, and bodyweight gain in an F<sub>2</sub> cross of  
292 Japanese quail using recursive models. *Poult. Sci.* 95:764-773. doi 10.3382/ps/pev376
- 293 Beck, P., M. Rodehutsord, J. Bennewitz, and W. Bessei. 2014. A pilot study of the genetic  
294 variation of phosphorus utilization in young Japanese quail (*Coturnix japonica*). *Poult.*  
295 *Sci.* 93:1916-1921. doi 10.3382/ps.2014-03982
- 296 Danzeisen, J. L., H. B. Kim, R. E. Isaacson, Z. J. Tu, and T. J. Johnson. 2011. Modulations of  
297 the chicken cecal microbiome and metagenome in response to anticoccidial and  
298 growth promoter treatment. *PLoS ONE* 6:e27949. doi 10.1371/journal.pone.0027949
- 299 Delezie, E., L. Maertens, and G. Huyghebaert. 2012. Consequences of phosphorus  
300 interactions with calcium, phytase, and cholecalciferol on zootechnical performance  
301 and mineral retention in broiler chickens. *Poult. Sci.* 91:2523-2531. doi  
302 10.3382/ps.2011-01937
- 303 de Verdal, H., A. Nary, D. Bastianelli, H. Chapuis, N. Mème, S. Urvoix, E. Le Bihan-Duval,  
304 and S. Mignon-Grasteau. 2011. *BMC Genetics* 12:71. doi:10.1186/1471-2156-12-71
- 305 Huber, K., E. Zeller, and M. Rodehutsord. 2015. Modulation of small intestinal phosphate  
306 transporter by dietary supplements of mineral phosphorus and phytase in broilers.  
307 *Poult. Sci.* 94:1009-1017. doi 10.3382/ps/pev065
- 308 Li, W., R. Angel, S.-W. Kim, K. Brady, S. Yu, and P. W. Plumstead. 2016. Impacts of dietary  
309 calcium, phytate, and nonphytate phosphorus concentrations in the presence or  
310 absence of phytase on inositol hexakisphosphate (IP6) degradation in different  
311 segments of broilers digestive tract. *Poult. Sci.* 95:581-589. doi 10.3382/ps/pev354

- 312 Maenz, D. D., and H. L. Classen. 1998. Phytase activity in the small intestinal brush border  
313 membrane of the chicken. *Poult. Sci.* 77:557-563.
- 314 Onyango, E. M., E. K. Asem, and O. Adeola. 2006. Dietary cholecalciferol and phosphorus  
315 influence intestinal mucosa phytase activity in broiler chicks. *Br. Poult. Sci.* 47:632-  
316 639.
- 317 Palacios, M. C., M. Haros, Y. Sanz, and C. M. Rosell. 2008. Selection of lactic acid bacteria  
318 with high phytate degrading activity for application in whole wheat breadmaking.  
319 *Food Sci. Techn.* 41:82-92. doi <http://dx.doi.org/10.1016/j.lwt.2007.02.005>
- 320 Ravindran, V., O. Adeola, M. Rodehutsord, H. Kluth, J. D. van der Klis, E. van Eerden, and  
321 A. Helmbrecht. Determination of ileal digestibility of amino acids in raw materials for  
322 broiler chickens – Results of collaborative studies and assay recommendations.  
323 Submitted to Poultry Science on 18 June, 2016
- 324 Rousseau, X., A.-S. Valable, M.-P. Létourneau-Montminy, N. Mème, E. Godet, M. Magnin,  
325 Y. Nys, M. J. Duclos, and A. Narcy. 2016. Adaptive response of broilers to dietary  
326 phosphorus and calcium restrictions. *Poult. Sci.* advance access. doi  
327 10.3382/ps/pew172
- 328 Schedle, K., R. Leitgeb, and M. Rodehutsord. 2016. Comparison of phosphorus digestibility  
329 between female and male broiler chickens. *Proc. Soc. Nutr. Physiol.* 25:56 (Abstr.)
- 330 Shastak, Y., and M. Rodehutsord. 2013. Determination and estimation of phosphorus  
331 availability in growing poultry and their historical development. *World's Poult. Sci. J.*  
332 69:569-585. doi 10.1017/S0043933913000585
- 333 Shastak, Y., and M. Rodehutsord. 2015. Recent developments in determination of available  
334 phosphorus in poultry. *J. Appl. Poult. Res.* 24:283-292. doi 10.3382/japr/pfv025
- 335 Taheri, H. R., H. Moravej, F. Tabandeh, M. Zaghari, and M. Shivazad. 2009. Screening of  
336 lactic acid bacteria toward their selection as a source of chicken probiotic. *Poult. Sci.*  
337 88:1586-1593. doi 10.3382/ps.2009-00041
- 338 Tamim, N. M., R. Angel, and M. Christman. 2004. Influence of dietary calcium and phytase  
339 on phytate phosphorus hydrolysis in broiler chickens. *Poult. Sci.* 83:1358-1367. doi  
340 10.1093/ps/83.8.1358
- 341 Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten. 2007.  
342 Handbuch der Landwirtschaftlichen Versuchs- und Untersuchungsmethodik  
343 (VDLUFA-Methodenbuch), Vol. III. Die chemische Untersuchung von Futtermitteln.  
344 VDLUFA-Verlag, Darmstadt, Germany.

- Witzig, M., A. Camarinha-Silva, R. Green-Engert, K. Hoelzle, E. Zeller, J. Seifert, L. E. Hoelzle, and M. Rodehutscord. 2015. Spatial variation of the gut microbiota in broiler chickens as affected by dietary available phosphorus and assessed by T-RFLP analysis and 454 pyrosequencing. *PLoS ONE* 10:e0143442. doi 10.1371/journal.pone.0143442
- WPSA (World's Poultry Science Association, Working Group 2: Nutrition of the European federation of branches). Determination of phosphorus availability in poultry. *World's Poult. Sci. J.* 69:687-698. doi 10.1017/S0043933913000688
- Yan, F., R. Angel, C. Ashwell, A. Mitchell, and M. Christman. 2005. Evaluation of the broiler's ability to adapt to an early moderate deficiency of phosphorus and calcium. *Poult. Sci.* 84:1232-1241. doi 10.1093/ps/84.8.1232
- Zeller, E., M. Schollenberger, I. Kühn, and M. Rodehutscord. 2015a. Effect of diets containing enzyme supplements and microwave-treated or untreated wheat on inositol phosphates in the small intestine of broilers. *Anim. Feed Sci. Techn.* 204:42-51. doi <http://dx.doi.org/10.1016/j.anifeedsci.2015.03.007>
- Zeller, E., M. Schollenberger, I. Kühn, and M. Rodehutscord. 2015b. Hydrolysis of phytate and formation of inositol phosphate isomers without or with supplemented phytases in different segments of the digestive tract of broilers. *J. Nutr. Sci.* 4:e1. doi 10.1017/jns.2014.62
- Zeller, E., M. Schollenberger, M. Witzig, Y. Shastak, I. Kühn, L. E. Hoelzle, and M. Rodehutscord. 2015c. Interactions between supplemented mineral phosphorus and phytase on phytate hydrolysis and inositol phosphates in the small intestine of broilers. *Poult. Sci.* 94:1018-1029. doi 10.3382/ps/pev087
- Zhang, W., S. E. Aggrey, G. M. Pesti, H. M. Edwards, Jr., and R. I. Bakalli. 2003. Genetics of phytate phosphorus bioavailability: heritability and genetic correlations with growth and feed utilization traits in a randombred chicken population. *Poult. Sci.* 82:1075-1079.



**Table 1.** Composition of the experimental diets.

Diet	A	B	C
<i>Ingredients (g/kg)</i>			
Soybean meal from de-hulled seed <sup>1</sup>	400.0	510.0	620.0
Corn starch	448.6	336.6	224.6
Limestone (finely ground)	7.4	9.4	11.4
Soybean oil	30.0	30.0	30.0
Dried egg albumen	18.0	18.0	18.0
Sucrose	80.0	80.0	80.0
Sodium chloride	3.0	3.0	3.0
DL-Methionine	2.7	2.7	2.7
L-Threonine	0.3	0.3	0.3
Titanium dioxide	5.0	5.0	5.0
Vitamin and trace element premix <sup>2</sup>	5.0	5.0	5.0
<i>Analyzed (g/kg, on dry matter basis)</i>			
CP	231	288	339
Total P <sup>3</sup>	3.02 (0.06)	3.80 (0.08)	4.59 (0.08)
Ca <sup>3</sup>	4.57 (0.09)	5.65 (0.09)	6.78 (0.13)
Ti <sup>3</sup>	3.20 (0.07)	3.18 (0.07)	3.17 (0.07)
InsP <sub>6</sub> -P	1.63	1.97	2.28
Ins(1,2,4,5,6)P <sub>5</sub> -P <sup>4</sup>	0.24	0.27	0.32

<sup>1</sup> The soybean meal contained per kg of dry matter (per analyses): 541 g crude protein, 76 g ash, 36 g crude fat, 41 g crude fiber, 116 g neutral detergent fiber, 72 g acid detergent fiber, 7.1 g P, 4.3 g InsP<sub>6</sub>-P, 3.3 g Ca.

<sup>2</sup> Premix provided the following (per kilogram of diet): vitamin A, 12,000 IU; cholecalciferol, 63 µg; vitamin E, 50 IU; vitamin K<sub>3</sub>, 1.5 mg; vitamin B<sub>1</sub>, 2.0 mg; vitamin B<sub>2</sub>, 7.5 mg; vitamin B<sub>6</sub>, 3.5 mg; vitamin B<sub>12</sub>, 20 mcg; niacin, 35 mg; D-pantothenic acid, 12 mg; choline chloride, 460 mg; folic acid, 1.0 mg; biotin, 0.2 mg; Fe, 80 mg (as FeSO<sub>4</sub>·H<sub>2</sub>O); Cu, 12 mg (as CuSO<sub>4</sub>·5H<sub>2</sub>O); Mn, 85 mg (as MnO); Zn, 60 mg (as ZnSO<sub>4</sub>·H<sub>2</sub>O); Co, 0.4 mg (as CoSO<sub>4</sub>·7H<sub>2</sub>O); I, 0.8 mg (as KI); Se, 0.15 mg (as Na<sub>2</sub>SeO<sub>3</sub>).

<sup>3</sup> Mean (standard deviation) of analyses made for each laboratory.

<sup>4</sup> All other lower inositol phosphate isomers were below the limit of quantification.

**Table 2.** Summary of the broiler chicken trials conducted in the collaborating stations.

Station number	Broiler strain	Sex	No of experimental replicates per diet	Birds per replicate	Age when placed on treatment diets (d)	Age at sampling (d)	BW before sampling (kg)	ADFI (g)	Feed/gain during the experimental period	Method of killing
1	Ross 708	male	8	8	20	25	1.12	123	1.63	CO <sub>2</sub> asphyxiation
2	Ross PM3	unsexed	8	8	23	28	1.86	174	1.22	Pentobarbital injection
3	Ross 308	unsexed	6 <sup>1</sup>	8	19	27	1.46	138	1.32	Cervical dislocation
4	Ross 308	unsexed	6	8	22	28	1.84	163	1.27	Cervical dislocation
5	Ross 308	male	6	10	22	27	1.03	81	1.26	CO <sub>2</sub> asphyxiation <sup>2</sup>
6	Ross 308	unsexed	6	8	23	28	1.52	151	1.06	Cervical dislocation
7	Ross 308	male	8	12	17	22	1.16	119	1.36	Cervical dislocation
8	Ross 308	male	6	12	20	27	1.57	133	1.35	Pentobarbital injection
9	Heritage 5632	male	6	8	16	21	0.82	93	1.43	CO <sub>2</sub> asphyxiation
10	Ross 708	unsexed	6	8	20	25	1.11	109	1.44	CO <sub>2</sub> asphyxiation
11	Ross 308	unsexed	7	8	18	23	1.03	93	1.42	CO <sub>2</sub> asphyxiation
12	Ross 308	male	8	8	16	25	1.45	98	1.32	Stunning
13	Ross PM3	unsexed	10	8	14	21	1.02	103	1.42	Cervical dislocation
14	Ross PM3	male	6 <sup>1</sup>	8	17	24	1.3	128	1.37	Pentobarbital injection
15	Ross 308	unsexed	6	10	18	28	1.72	140	1.40	Cervical dislocation
16	Ross PM3	unsexed	10 <sup>1</sup>	4	12	22	1.00	70	1.28	CO <sub>2</sub> asphyxiation
17	Ross 308	male	6	12	14	24	1.24	96	1.31	Sedation and Ketamin

<sup>1</sup>Each replicate comprised two pooled cages.<sup>2</sup>CO<sub>2</sub> asphyxiation following anesthesia with a gas mixture of 35% CO<sub>2</sub>, 35% N<sub>2</sub>, and 30% O<sub>2</sub>.

**Table 3.** Prececal digestibility of P and Ca of diets with variable inclusion of soybean meal and limestone as determined in 16 trials with broiler chickens

Diet	No of station	n <sup>2</sup>	P digestibility <sup>1</sup> (%)						Ca digestibility <sup>1</sup> (%)					
			A		B		C		A		B		C	
			Mean	SD	Mean	sd	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	1	8	58	2.7	52	5.0	48	4.9	47	8.9	47	6.6	39	8.6
2	2	8 <sup>3</sup>	67	3.2	58	3.3	52	1.2	64	3.4	55	5.6	47	3.3
3	3	6	68	4.8	62	2.2	50	3.2	56	5.0	49	3.1	42	3.4
4	4	6 <sup>4</sup>	63	3.9	56	3.7	52	2.1	65	9.2	49	4.8	44	3.5
5	5	6	73	1.9	69	2.9	66	2.9	62	2.5	58	2.3	54	3.8
6	6	6	61	3.8	60	4.1	53	7.2	55	7.5	51	11.6	41	7.4
7	7	8	59	5.3	54	3.2	50	4.0	50	4.9	44	4.8	39	3.4
8	8	6	71	2.2	62	4.9	62	3.6	62	2.9	53	5.5	54	4.2
9	9	6	62	4.4	55	2.9	48	3.9	55	7.5	45	5.6	39	6.6
10	10	6	68	2.5	61	2.9	53	1.5	58	4.0	52	6.5	43	3.7
11	11	7	55	5.5	46	5.3	45	6.2	56	6.3	46	5.1	43	5.7
12	12	8	71	4.0	61	4.2	55	5.2	64	4.3	55	3.8	49	4.6
13	13	10	72	5.6	70	6.3	62	2.7	58	5.8	55	9.7	51	6.4
15	15	6	74	2.3	63	9.7	61	6.9	62	1.7	51	8.7	50	7.8
16	16	10	72	2.8	66	4.9	59	4.9	44	5.7	45	7.6	43	7.7
17	17	6	82	1.6	79	3.2	69	2.4	71	3.4	68	2.7	58	4.0

Diets: A = 400 g/kg soybean meal; B = 525 g/kg soybean meal; C = 650 g/kg soybean meal.

<sup>1</sup>Effects of diet, station, and their interaction were statistically significant ( $P < 0.05$ ).

<sup>2</sup>Number of replicates (cages) per diet. <sup>3</sup>n = 7 for diet C. <sup>4</sup>n = 5 for diets A and C.

**Table 4.** Results of linear regression analysis of prececal digestible P concentration (y, g/kg DM) in function of dietary P concentration (x, g/kg DM) using diets with incremental inclusion levels of soybean meal

No of station	Intercept	<i>SE</i>	Slope		<i>SE</i>	R <sup>2</sup>	Root MS error
1	0.83	<i>0.30</i>	0.31	<sup>abc</sup>	<i>0.07</i>	>0.99	0.01
2	1.37	<i>0.41</i>	0.22	<sup>bc</sup>	<i>0.08</i>	0.97	0.02
3	1.50	<i>0.43</i>	0.19	<sup>c</sup>	<i>0.07</i>	0.68	0.09
4	0.99	<i>0.40</i>	0.30	<sup>abc</sup>	<i>0.07</i>	0.98	0.03
5	0.66	<i>0.42</i>	0.51	<sup>a</sup>	<i>0.08</i>	>0.99	<0.01
6	0.72	<i>0.42</i>	0.38	<sup>abc</sup>	<i>0.08</i>	0.93	0.07
7	0.79	<i>0.42</i>	0.33	<sup>abc</sup>	<i>0.08</i>	>0.99	<0.01
8	0.80	<i>0.42</i>	0.43	<sup>ab</sup>	<i>0.07</i>	0.95	0.06
9	1.34	<i>0.42</i>	0.19	<sup>c</sup>	<i>0.08</i>	0.91	0.04
10	1.36	<i>0.43</i>	0.25	<sup>bc</sup>	<i>0.07</i>	0.94	0.04
11	0.81	<i>0.42</i>	0.27	<sup>bc</sup>	<i>0.08</i>	0.92	0.05
12	1.34	<i>0.42</i>	0.26	<sup>bc</sup>	<i>0.07</i>	>0.99	0.02
13	0.94	<i>0.42</i>	0.42	<sup>ab</sup>	<i>0.07</i>	0.95	0.07
15	1.02	<i>0.41</i>	0.38	<sup>abc</sup>	<i>0.07</i>	0.97	0.04
16	1.12	<i>0.41</i>	0.34	<sup>abc</sup>	<i>0.08</i>	>0.99	0.02
17	1.37	<i>0.42</i>	0.41	<sup>abc</sup>	<i>0.08</i>	0.91	0.08

Superscript letters indicate significant differences between slopes.

**Table 5.** Prececal disappearance of InsP<sub>6</sub> of diets with variable inclusion of soybean meal as determined in 4 trials with broiler chicken


Diet	No of station	n <sup>2</sup>	Prececal InsP <sub>6</sub> disappearance <sup>1</sup> (%)					
			A		B		C	
			Mean	SD	Mean	SD	Mean	SD
4		6 <sup>3</sup>	70	4.5	52	8.5	47	5.0
5		6	74	3.8	67	4.7	58	7.1
7		8	58	10.8	47	8.1	36	8.9
9		6	62	4.3	43	3.6	23	11.4

Diets: A = 400 g/kg soybean meal; B = 525 g/kg soybean meal; C = 650 g/kg soybean meal.


<sup>1</sup>Effects of diet, station, and their interaction were statistically significant ( $P < 0.05$ ).

<sup>2</sup>Number of replicate cages.


<sup>3</sup>n = 5 for diets A and C.




**Figure 1.** Scatter plot of prececal digestibility data from 16 broiler chicken trials conducted in 16 stations and using diets containing 400 (A), 525 (B) or 650 (C) g/kg soybean meal (each dot shows the mean of one station for the respective diet).



**Figure 2.** Relationship between the concentration of InsP<sub>6</sub>-P in the diet and prececal InsP<sub>6</sub>-P disappearance in diets with different levels of soybean meal and studied in 4 broiler trials in 4 stations (means and SD). Slopes estimated for stations 4, 5, and 7 did not deviate from zero ( $P > 0.05$ ).




**Figure 3.** Relationship between the concentration of prececal digestible P in dependence of prececal InsP<sub>6</sub>-P disappearance in diets including different levels of soybean meal and studied in 4 broiler trials in 4 stations (each symbol shows the value of one cage). The slope of the linear regressions calculated for each diet were not significantly different ( $P = 0.27$ ), and the pooled slope was 0.91.




**Figure 4.** Relationship between the concentration of prececal digestible P and prececal digestible Ca in diets fed to broilers and using diets containing 400 (A), 525 (B) or 650 (C) g/kg soybean meal (each symbol shows the mean of one out of 16 stations for the respective diet). The slopes determined separately for each diet were not significantly different.


Prececal P digestibility of diet A (%)




Prececal P digestibility of diet A (%)




Prececal P digestibility of diet A (%)




Prececal P digestibility of diet A (%)




Prececal P digestibility of diet A (%)



Prececal P digestibility of diet A (%)



Prececal P digestibility of diet A (%)





**Figure 5.** Comparison between the prececal digestibility of P in diet A with the formulated P concentration of the pre-experimental starter feed (panel A), the supplemented phytase of the pre-experimental starter feed (panel B), the age of birds when the experimental diets were introduced (panel C), the duration of the experimental feeding period (panel D), the age of birds at the day of sampling (panel E), the BW of the birds at the day of sampling (panel F), and the ADFI in the trial period (panel G). Each dot shows the mean of one station. The difference to zero of the calculated slopes of linear regressions was significant in panel D ( $P < 0.01$ ), but not in panel A ( $P = 0.08$ ) or any other panel ( $P > 0.1$ ).